Virginia Racing Commission Medication Synopsis – January 2017

- 1. Furosemide is the only medication permitted within the 24-hour period prior to post and must be administered IV only to listed horses, no later than 4 hours prior to post (3 hours at designated ship-in meets only) on the grounds of the racetrack, by the Lasix veterinarian;
- 2. The minimum dosage of furosemide is 3 ml (2 ml at designated ship-in meets);
- 3. Private veterinarians are prohibited from contact with a horse on race day, prior to post;

30 controlled therapeutic medications, with laboratory thresholds and withdrawal guidelines are listed below (note: the recommendation for methylprednisolone includes a strong caution to reflect more recent experience):

<u>Drug</u>	<u>Dose/Route</u>	<u>Withdrawal</u>	<u>Threshold</u>	
Acepromazine	0.05 mg/kg IV	48 hours	10 ng/ml urine	
Albuterol	720 mcg Intra-Nasal qid	72 hours	1 ng/ml urine	
Betamethasone (IA only)	9 mg – 1 joint	7 days	10 pg/ml plasma	
Butorphanol	0.1 mg/kg IV	48 hours	2 ng/ml plasma	
Cetirizine*	0.4 mg/kg bid x 5days	48 hours*	6 ng/ml plasma	
*Do not administer ive	ermectin w/I 24 hours if using cet	tirizine		
Cimetidine	20 mg/kg bid 7 doses	24 hours	400 ng/ml plasma	
Clenbuterol	0.8 mcg/kg PO bid	14 days	LOD plasma	
Dantrolene	500 mg PO	48 hours	100 pg/ml plasma	
Detomidine	5 mg IV (once)	48 hours	1 ng/ml plasma	
Dexamethasone	0.05 mg/kg IV, IM, PO	72 hours	5 pg/ml plasma	
Diclofenac (topical)	5" ribbon topically	48 hours	5 ng/ml plasma	
DMSO	50 ml IV	48 hours	10 mcg/ml plasma	
Firocoxib	0.1 mg/kg PO x 4d	14 days	20 ng/ml plasma	
Flunixin	1.1 mg/kg IV (once)	32 hours	20 ng/ml plasma	
Furosemide	150 - 500 mg IV	3 hours	100 ng/ml plasma	
Glycopyrrolate	1 mg total IV	48 hours	3 pg/ml plasma	
Guaifenesin	2 gm PO bid 5 doses	48 hours	12 ng/ml plasma	
Isoflupredone	10 mg SQ; 20 mg IA 1 joint	7 days	100 pg/ml plasma	
Ketoprofen	2.2 mg/kg IV	24 hours	2 ng/ml plasma	
Lidocaine	200 mg total SQ	72 hours	20 pg/ml plasma	
Mepivacaine	0.07 mg/kg SQ	72 hours	LOD plasma	
Methocarbamol	15 mg/kg IV or 5 g PO	48 hours	1 ng/ml plasma	
Methylprednisolone** (IA)	100 mg – 1 joint	21 days**	100 pg/ml plasma	
**Methylprednisolone should not be used or if used the trainer should have the horse tested at				

**Methylprednisolone should not be used, or if used the trainer should have the horse tested at his expense prior to entry to ensure the serum concentration is below the regulatory limit.

Omeprazole	2.2 g PO x 4 days	24 hours	10 ng/ml plasma
Phenylbutazone (IV only)	4.0 mg/kg IV	24 hours	2 mcg/ml plasma
Prednisolone	1 mg/kg PO	48 hours	1 ng/ml plasma
Procaine Penicillin***	IM	30 days***	25 ng/ml plasma

***Use within 30 days must be reported to the racing commission, and the horse must be kept under security guard surveillance/quarantine for 6 hours prior to post, at trainer's expense.

Ranitidine	8 mg/kg bid x 7 doses	24 hours	40 ng/ml plasma
Triamcinolone Acetonide (IA)	9 mg – IA 1 joint	7 days	100 pg/ml plasma
Xylazine	200 mg IV	48 hours	200 pg/ml plasma
			LOD = level of detection

Cobalt Regulation:

The RMTC tiered penalty structure for Cobalt:

- 1. For horses testing above 25 nanograms of cobalt per mL of plasma, but below 50 nanograms of cobalt per mL of plasma:
 - -- First offense trainer will receive a fine or a warning
 - -- Horse will remain on the Veterinarian's List until it tests below 25 nanograms/mL (This testing is performed at owner/trainer's expense)
 - 2. For horses testing above 50 nanograms of cobalt per mL of plasma:
 - -- Class B penalty (RCI Model Rule for a first offense)
 - --fine (\$500-\$1,000)
 - --suspension (15-60 days)
 - --loss of purse
 - --- Horse will remain on the Veterinarian's List until it tests below 25 nanograms/mL (This testing is at owner/trainer's expense)

For more information about medications, go to the Thoroughbred Horsemen's Association website: www.tharacing.com